

Colonoscopy

CLENPIQ- AM Procedure Instructions

Procedure Date: _____ Check-in Time: _____ GI Physician: _____

Location: _____ Procedure Time: _____

Special Instructions:

To ensure a successful exam, please follow all instructions on this sheet carefully. Failure to do so could result in your procedure being postponed or cancelled.

post posed or canceled.

If you are pregnant, please discuss the risks and benefits of this procedure with your physician.

If you take anti-coagulants / blood thinners (for example: Coumadin, Plavix, Pradaxa, Xarelto, Eliquis, Effient, Brilinta, etc.) or Insulin, please notify our office at least 10 days prior to your procedure.

Please inform us if you weigh greater than 250 lbs., have kidney disease, congestive heart failure, artificial heart valves, a pacemaker or defibrillator (AICD), sleep apnea, or ascites.

Please discontinue iron or fiber supplements, seeds, nuts, popcorn, and raw vegetables 3 days before your colonoscopy.

Transportation:

- Please make arrangements to have someone drive you to and from the designated location for your procedure.
- You will not be allowed to use any form of public transportation following your procedure, which includes: Taxi, Bus, Ride-share service, Uber, Lyft and the like.
- The test may be cancelled if you do not have a driver, there will be no exceptions.
- You will be under sedation during your procedure and will be unable to work, drive or operate heavy machinery the day of your procedure.

Pick Up Your Prescription:

You may fill your **CLENPIQ** prescription at any pharmacy. IT IS VERY IMPORTANT THAT YOU REFER TO THESE PREP INSTRUCTIONS FROM YOUR DOCTOR REGARDING WHEN YOU NEED TO STOP EATING AND DRINKING. **DO NOT FOLLOW THE INSTRUCTIONS THAT YOU RECEIVE FROM THE PHARMACY OR WHAT IS ON THE PREP PACKAGE.**

The Day Before Procedure:

The day before your colonoscopy is when you will start your preparation. The colon needs to be as clean of stool as possible. If this cannot be achieved, the test may be incomplete, and abnormalities may be missed.

- Do NOT use any tobacco products or marijuana, do NOT take non-prescribed drugs, and do NOT drink alcohol after midnight before your procedure.
- DRINK ONLY CLEAR LIQUIDS ALL DAY (No solid food):

It is important to prepare your body with plenty of clear liquids before the prep. The explanation of a clear liquid diet is anything you can see through if held up to a light.

Approved Clear Liquids:

- Water, Black Coffee or Black Tea (no cream or artificial cream)
 - Strained fruit juices WITHOUT pulp (Example: apple, white grape)
 - Clear Broth or Bouillon (Example: chicken or beef or vegetable broth)
 - Carbonated CLEAR soft drinks (Example: Sprite, ginger ale)
 - Gatorade, Kool aid or lemonade with no pulp (NO RED OR PURPLE)
 - Plain Jell-O (without fruit or toppings) (NO RED OR PURPLE)
 - Popsicles (NO RED OR PURPLE)
 - NO DAIRY PRODUCTS
 - NO ALCOHOLIC BEVERAGES
- Taking The Prep: First Dose of CLENPIQ
Starting At 12PM Noon (Day before Procedure)
Step 1: Take (2) Dulcolax tablets with 8oz of water.
At 4PM
Step 2: Drink all of one bottle(160mL) of CLENPIQ. You may prefer to drink the liquid through a straw.
Step 3: You must drink at least five more 8oz glasses of water/clear liquid over the next five hours. Very Important.



- Taking The Prep: Second Dose of CLENPIQ

Starting at 10PM

REPEAT STEPS 2 AND 3, drinking the entire prep solution (VERY IMPORTANT) and at least 5 more glasses of clear liquid to follow. You must finish drinking all liquids at least 3 hours before the procedure time.

-Bowel movements usually start within 1 or 2 hours after you begin drinking and usually continue for 1 or 2 hours after you finish.

-It is normal to feel full and somewhat bloated while you are drinking the prep.

If you become nauseated, slow your pace of drinking.

Day Of Your Procedure:

- In the morning, you should use inhalers, and take heart, blood pressure, seizure, anti-anxiety, and respiratory medications with a sip of water, enough to swallow them (no blood thinners or oral diabetic medications). You should wake up early and take these at least 2 hours before your test, even if you are having an EGD with your colonoscopy.
- On the morning of the procedure, you may pass some cloudy or dark liquid stool. This is common and not something to be concerned about. If you drank the prep until your stool was clear the day before, your prep is probably fine.
- Female patients of childbearing age will be required to provide a urine specimen for a pregnancy test prior to anesthesia the day of the procedure. Please avoid urinating prior to arrival the day of your procedure to prevent delay or possible cancellation.

Questions:

- If you have questions, or if you find you cannot tolerate the prep, then please call our office at 601-450-5162

